

2.03.01

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	1 of 18

Last Review Status/Date: June 2015

In Vitro Chemoresistance and Chemosensitivity Assays

Description

In vitro chemoresistance and chemosensitivity assays have been developed to provide information about the characteristics of an individual patient's malignancy to predict potential responsiveness of their cancer to specific drugs. Thus, these assays are sometimes used by oncologists to select treatment regimens for an individual patient. Several assays have been developed that differ with respect to processing of biological samples and detection methods. However, all involve similar principles and share protocol components including: 1) isolation of cells and establishment in an in vitro medium (sometimes in soft agar); 2) incubation of the cells with various drugs; 3) assessment of cell survival; and 4) interpretation of the result.

Background

A variety of chemosensitivity and chemoresistance assays have been clinically evaluated in human trials. All assays use characteristics of cell physiology to distinguish between viable and non-viable cells to quantify cell kill following exposure to a drug of interest. With few exceptions, drug doses used in the assays are highly variable depending on tumor type and drug class, but all assays require drug exposures ranging from several-fold below physiologic relevance to several-fold above physiologic relevance. Although a variety of assays exist to examine chemosensitivity or chemoresistance, only a few are commercially available. Available assays are outlined as follows:

Methods using differential staining/dye exclusion:

- The Differential Staining Cytotoxicity Assay (1) This assay relies on dye exclusion of live cells after mechanical disaggregation of cells from surgical or biopsy specimens by centrifugation. Cells are then established in culture and treated with the drugs of interest at three dose levels; the middle dose is that which could be achieved in therapy; 10-fold lower than the physiologically relevant dose; and, 10-fold higher. Exposure time ranges from 4-6 days; then, cells are restained with fast green dye and counterstained with hematoxylin and eosin (H&E). The fast green dye is taken up by dead cells, and H&E can differentiate tumor cells from normal cells. The intact cell membrane of a live cell precludes staining with the green dye.

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	2 of 19

Drug sensitivity is measured by the ratio of live cells in the treated samples to the number of live cells in the untreated controls.

- The EVA/PCD™ assay (available from Rational Therapeutics, Long Beach, CA). This assay relies on ex-vivo analysis of programmed cell death, as measured by differential staining of cells after apoptotic and non-apoptotic cell death markers in tumor samples exposed to chemotherapeutic agents. Tumor specimens obtained through biopsy or surgical resection is disaggregated using DNase and collagenase IV to yield tumor clusters of the desired size (50-100 cell spheroids). Because these cells are not proliferated, these micro-aggregates are believed to more closely approximate the human tumor micro-environment. These cellular aggregates are treated with the dilutions of the chemotherapeutic drugs of interest and incubated for 3 days. After drug exposure is completed, a mixture of Nigrosin B & Fast Green dye with glutaraldehyde-fixed avian erythrocytes added to the cellular suspensions. (2) The samples are then agitated and cytospin-centrifuged and, after air drying, are counter-stained with H&E. The endpoint of interest for this assay is cell death as assessed by observing the number of cells differentially stained due to changes in cellular membrane integrity. (3)
- The fluorometric microculture cytotoxicity assay (FMCA) is another cell viability assay that relies on the measurement of fluorescence generated from cellular hydrolysis of fluorescein diacetate to fluorescein in viable cells. (4) Cells from tumor specimens are incubated with cytotoxic drugs; drug resistance is associated with higher levels of fluorescence.

Methods using incorporation of radioactive precursors by macromolecules in viable cells:

- Tritiated thymine incorporation measures uptake of tritiated thymidine by DNA of viable cells. Using proteases and DNase to disaggregate the tissue, samples are seeded into single cell suspension cultures on soft agar. They are then treated with the drug(s) of interest for 4 days. After 3 days, tritiated thymidine is added. After 24 hours of additional incubation, cells are lysed and radioactivity is quantified and compared to a blank control consisting of cells that were treated with sodium azide. Only cells that are viable and proliferating will take up the radioactive thymidine. Therefore, there is an inverse relationship between uptake of radioactivity and sensitivity of the cells to the agent(s) of interest. (5)
- The Extreme Drug Resistance assay (EDR®) (6) (commercially available at Exiqon Diagnostics, Tustin, CA) is methodologically similar to the thymidine incorporation assay, using metabolic incorporation of tritiated thymidine to measure cell viability; however, single cell suspensions are not required, so the assay is simpler to perform. Tritiated thymidine is added to the culture and uptake is quantified after various incubation times. Only live (resistant) cells will incorporate the compound. Therefore, the level of tritiated thymidine incorporation is directly related to chemoresistance. The interpretation of the results is unique in that resistance to the drugs is evaluated as opposed to evaluation of responsiveness. Tumors are considered to be highly resistant when thymidine incorporation is at least 1 standard deviation (SD) above reference samples

Methods to quantify cell viability by colorimetric assay:

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	3 of 19

- The Histoculture Drug Resistance Assay (HDRA; AntiCancer Inc., San Diego, CA). (7) This assay evaluates cell growth after chemotherapy treatment based on a colorimetric assay that relies on mitochondrial dehydrogenases in living cells. Drug sensitivity is evaluated by quantification of cell growth in the 3-dimensional collagen matrix. There is an inverse relationship between the drug sensitivity of the tumor and cell growth. Concentrations of drug and incubation times are not standardized and vary depending on drug combination and tumor type.

Methods using incorporation of chemoluminescent precursors by macromolecules in viable cells:

- The Adenosine Triphosphate (ATP) Bioluminescence Assay. This assay relies on measurement of ATP to quantify the number of viable cells in a culture. Single cells or small aggregates are cultured, and then exposed to drugs. Following incubation with drug, the cells are lysed and the cytoplasmic components are solubilized under conditions that will not allow enzymatic metabolism of ATP. Luciferin and firefly luciferase are added to the cell lysis product. This catalyzes the conversion of ATP to adenosine di- and monophosphate and light is emitted proportionally to metabolic activity. This is quantified with a luminometer. From the measurement of light, the number of cells can be calculated. A decrease in ATP indicates drug sensitivity, whereas no loss of ATP suggests that the tumor is resistant to the agent of interest.
- ChemoFX® (Helomics Corporation, previously called Precision Therapeutics, Pittsburgh, PA). (8) This assay also relies on quantifying ATP based on chemoluminescence. Cells must be grown in a monolayer rather than in a 3-dimensional matrix.

Methods using differential optical density

- CorrectChemo® (previously called the Microculture Kinetic [MiCK] assay) (Diatech Oncology, Franklin, TN). (9) Similar to the EVA/PCD assay, this assay relies on measures of programmed cell death. In the assay, tumor cells are exposed to multiple concentrations of drugs and cultured. The optical density of the cells is measured over time, to create a density-by-time curve. A sudden increase in optical density is associated with cell apoptosis; the extent of drug-induced apoptosis is a measure of the cell's sensitivity to that agent.

The rationale for chemosensitivity assays is strongest where there are a variety of therapeutic options and there are no clear selection criteria for any particular regimen in an individual patient.

Regulatory Status

Commercially available chemosensitivity and chemoresistance assays are laboratory developed tests for which approval from the U.S. Food and Drug Administration (FDA) is not required when the tests are performed in a laboratory licensed by the Clinical Laboratory Improvement Act (CLIA) for high-complexity testing. Such tests must meet the general regulatory standards of the Clinical Laboratory Improvement Act (CLIA).

Related Policies

None

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	4 of 19

Policy

**This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.*

In vitro chemosensitivity assays, including, but not limited to, the Histoculture Drug Response Assay, a fluorescent cytoprint assay, the ChemoFx assay, or the CorrectChemo assay, are considered **not medically necessary**.

In vitro chemoresistance assays, including, but not limited to, extreme drug resistance assays, are considered **not medically necessary**.

Rationale

This policy is based on a 2002 Technology Evaluation Center (TEC) Assessment (10), and a 2004 *Journal of Clinical Oncology* systematic review, (11) which concluded that evidence is insufficient to support use of chemosensitivity and chemoresistance assays for guiding choice of therapy regimen in cancer patients.

A variety of studies have reported a correlation between in vitro prediction or response and clinical response. While these studies may have internal validity, they cannot answer the question of whether patients given assay-guided therapy or empiric therapy have different outcomes. To determine whether assay-guided treatment results in overall different outcomes than empiric treatment, it is important to take into account response rates, survival, adverse effects and quality of life. These effects may be assessed indirectly, for example, using decision analysis, or directly with comparative trials. Both the 2002 BCBSA TEC Assessment and the 2004 systematic review (10, 11) recommend validating chemotherapy sensitivity and resistance assays with direct evidence gathered from prospective trials comparing patients treated empirically to patients treated with assay-directed therapy. In this way, not only can response rates and survival be taken into account, but also adverse events (eg, from the toxic effects of an ineffective drug or delay or loss of benefits of an effective drug) and quality of life.

Chemoresistance Assays

Chemoresistance assays are used to deselect potential chemotherapeutic regimens. The negative predictive value (NPV) is a key statistical measure. Unless the NPV is high, there is a chance that clinical decision making based on a chemoresistance assay could inappropriately exclude an effective therapy. The NPV will vary according to the prior probability of chemoresistance, as well as the assay's sensitivity and specificity. The 2002 TEC Assessment (10) concluded that chemoresistance assays have the highest clinical relevance in tumors with low probability of response. The EDR assay was specifically designed to produce a very high NPV (>99%), such that the possibility of inappropriately excluding effective chemotherapy is remote in all clinical situations. (12)

To determine whether chemoresistance assays have value in clinical decision making, studies comparing outcomes for patients managed with chemoresistance assays to those managed with

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	5 of 19

routine care would be ideal. Potential relevant clinical outcomes include improved survival and avoidance of toxicity (as an intermediate outcome).

The bulk of the literature regarding EDR assays have focused on correlational studies that correlate results from predictive in vitro assays with observed outcomes of chemotherapy. However, in these studies, the patients do not receive assay-guided chemotherapy regimens. As discussed in the 2004 systematic review, (11) correlational studies are inadequate for several reasons. First, such studies often aggregate patients with different tumor types, disease characteristics, chemotherapy options, and probabilities of response. This process is problematic, since the accuracy of each assay used to predict in vivo response probably varies across different malignancies and patient characteristics.

Second, the method by which assay results are translated into treatment decisions is not standardized. Third, it is important to consider not only response but also survival, quality of life, and adverse effects. The overall value of assay-guided therapy depends on the net balance of all health outcomes observed after treatment for all patients subjected to testing, regardless of the assay results or the accuracy of its predication for response. Examples of some of the earlier published correlation studies of the EDR assay include those by Eltabbakh et al (13, 14), Mehta et al, (15) Holloway et al, (16) and Ellis et al. (17)

The 2002 TEC Assessment identified 1 nonrandomized retrospective comparative study using the extreme drug resistance (EDR®) assay, published by Loizzi et al. in 2003. (18) While this study of patients with recurrent ovarian cancer found a significantly higher overall response rate, better progression-free survival (PFS), and higher overall survival (OS) among platinum-sensitive patients receiving assay-guided therapy, it was not designed to adequately address potential biases and confounding. Since the Loizzi et al paper appeared, no additional comparative studies of assay-guided therapy versus physician-directed therapy have appeared for chemoresistance assays.

Correlational Studies

Prospective. A study by Tiersten et al. (19) was designed to use the Oncotech EDR assay to examine whether chemotherapy resistance was an independent predictor of PFS in patients with ovarian cancer treated with neoadjuvant chemotherapy and surgical cytoreduction followed by intraperitoneal chemotherapy. Fifty-eight eligible women were prospectively enrolled in this study; however, results from the EDR assay were not used to direct therapy. Evaluable EDR assay results were available for 22 of the 58 patients. No difference in PFS was reported. Follow-up has not been sufficient to measure OS. These data do not provide support for use of the EDR assay in predicting outcome and guiding patient management.

A 2006 review published by Nagoury et al included 21 non-comparative studies using ex-vivo programmed cell death assays. The authors of these studies correlated the drug susceptibility findings of the ex-vivo assay with objective clinical response (complete or partial) compared to non-responders for 659 total patients. The authors obtained aggregate positive values by site of primary cancer: breast (82.9%), colon (80%), non-small-cell lung cancer (66.7%), gynecologic (77%), and small-cell lung cancer (50%).(3). A 2012 study by this same investigator prospectively assessed 98 patients with non-small-cell lung cancer treated between 2003 and 2010.(2) Only 41 were found to be eligible for

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	6 of 19

inclusion and were tested with the EVA/PCD™ assay to determine which chemotherapeutic drugs to use. A further 10 patients were excluded (5 due to insufficient cellular yield, 3 for resistance to all drugs tested, and 2 due to physician's choice) yielding only 31 patients who received the assay-recommended treatment. The authors compared the results of these 31 patients treated with assay-directed chemotherapy to historic controls (not described) on the outcome of observed objective response rate (complete response and partial response). The objective response rate for the study was 64.5% (95% confidence interval [CI], 46.9 to 78.9%) which was significantly greater than the stated historic standard of 30% objective response ($p < 0.001$).

Retrospective. In 2010, Matsuo et al published a study examining the relevance of EDR in epithelial ovarian carcinomas. (20) Two-hundred fifty-three records from the Oncotech database were identified for women with advanced stage ovarian cancer and from whom samples were collected at the time of the primary surgery. Tissue samples were cultured and tested for response to primary drugs (4 platinum- or taxane-based) and secondary drugs (eg, gemcitabine, topotecan, doxorubicin, etoposide, 5-fluorouracil (5-FU). Paclitaxel showed the highest resistance rate. Other agents had a resistance rate of less than 20%. There was only one (0.4%) tumor that showed complete resistance to all drugs tested; and 25% of tumors showed no resistance to any of the drugs. There was no statistical correlation between assay results and response to initial chemotherapy. The investigator acknowledges that the study, due to its retrospective and noncomparative design is not sufficiently strong to validate use of this assay in managing therapy. Potential confounding factors, as described by the investigator, may have included tumor heterogeneity and the variations in resistance between primary tumor and metastases.

Another study by the same group (21) evaluated the role of the EDR assay to platinum- and taxane-based therapies for management of advanced epithelial ovarian, fallopian, and peritoneal cancers. From the Oncotech database, 173 cases were identified. For all cases, tissue was collected at the time of cytoreductive therapy. The EDR assay was performed on all samples, and tumors were classified as having low drug resistance (LDR), intermediate drug resistance (IDR), or extreme drug resistance (EDR). The 58 patients (33.5%) whose tumors had LDR to both platinum and taxane showed statistically improved PFS and OS compared to the 115 patients (66.5%) who demonstrated IDR or EDR to platinum and/or taxane (5-year OS rates, 41.1% vs 30.9%, respectively; $p = 0.014$). The 5-year OS rates for the 28 (16.2%) cases that had optimal cytoreduction with LDR to both platinum and taxane was significantly improved over the 62 (35.8%) cases that were suboptimally cytoreduced with IDR or EDR to platinum and/or taxane (54.1% vs 20.4%, respectively; $p < 0.001$). Although the EDR assay was predictive for survival, it is of interest that assay results did not indicate response to therapy with either taxane or cisplatin. The investigators conclude that the EDR assay may be an independent predictor of PFS and OS; however, a prospective, randomized trial would be required to further assess its clinical utility in predicting response to taxane or platinum therapies.

A smaller study by Matsuo et al testing the EDR assay for prediction of uterine carcinosarcoma response to taxane and platinum was also conducted. (22) Of 51 cases, 31 (60.8%) received postoperative chemotherapy with at least a single agent; and 17 (33.3%) received combination chemotherapy with platinum and taxane modalities. Overall response rate for the 17 combination chemotherapy cases was 70.6%. Presence of EDR to either platinum or taxane showed a significantly lower PFS (1-year PFS rate, 28.6% vs. 100%, respectively; $p = 0.01$) and lower OS (5-year OS rate,

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	7 of 19

26.9% vs. 57.1%, respectively; $p=0.033$). These data indicate that use of an in vitro drug resistance assay may be predictive of response to chemotherapy response and survival outcome in advanced ovarian and uterine carcinosarcoma. However, larger, prospective, randomized clinical trials (RCTs) would be required to validate use of this assay for directing chemotherapy regimens.

Matsuo et al also completed a study examining the rates of EDR after cytoreductive therapy and neoadjuvant chemotherapy versus the rates of ERD after postoperative chemotherapy. (23) The goal of this study was not to test whether the EDR assay could direct therapeutic regimens. The findings suggested that platinum resistance was most common after neoadjuvant chemotherapy, while paclitaxel resistance was more prevalent after postoperative chemotherapy.

Karam et al conducted a retrospective review of 377 patients with epithelial ovarian cancer to examine the effect of EDR assay-guided therapy on outcomes in the primary and recurrent setting. (24) The primary endpoints were time to progression (TTP), OS, and survival after recurrence (RS). The patient population was heterogeneous, with a median age of 59 years (median 24-89), tumor completely resected in 30% of patients and varying tumor stages (Federation of Gynecologists and Obstetricians [FIGO] stages I, II, III, and IV in 7%, 4%, 78%, and 11%, respectively). Sixty-four percent of patients underwent a secondary cytoreductive surgery. Patients had an EDR assay sent either at the time of their primary cytoreductive surgery ($n=217$) or at the time of disease recurrence ($n=160$). Predictors of survival included increasing age and greater volume of residual disease after cytoreductive surgery. EDR assay results analyzed for single agents or combinations of chemotherapies failed to independently predict patient outcomes regardless of whether the assay was performed at the time of the primary surgery or at recurrence.

Hetland et al conducted a study to identify primary platinum resistance in epithelial ovarian cancer patients with FIGO stage III-IV disease. (25) Eighty-five biopsies from 58 patients were included in the study. Resistance was assessed with a modified drug-response assay including ATP-based tumor-chemosensitivity and EDR assay. Samples were tested for response to platinum, paclitaxel and the combination of the drugs. Results from the assay were combined, and tumors were classified using a resistance index, which summarized the percentage of tumor growth inhibition for each drug concentration tested. All patients received a primary chemotherapy treatment of carboplatin, paclitaxel or a combination of the two drugs. Platinum resistance, as defined by the risk index, was associated with significantly poorer PFS ($p=0.03$) with a median value of 3.9 months (95% CI, 3.2 to 4.7) compared with the platinum sensitive group with a median PFS of 8.1 months (95% CI, 3.7 to 12.4). Patients who had partial response, stable disease or progressive disease were more resistant to platinum based on risk index score than those with a complete response ($p=0.02$). In a sub-group analysis of metastatic tumors, platinum resistance was not associated with PFS or clinical response. Response to paclitaxel or carboplatin/paclitaxel was not associated with PFS or clinical response. In vitro response was not associated with overall survival in any group.

Comparative Studies Testing Outcome with Assay-Directed Therapy Versus Physician-Chosen Therapy

None identified

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	8 of 19

Section Summary:

Some retrospective and prospective studies suggest that chemoresistance assays, particularly the EDR assay, may be associated with chemotherapy response. However, prospective studies do not consistently demonstrate that chemoresistance assay results are associated with survival. Furthermore, no comparative studies were identified that compare outcomes between patients managed with assay-directed therapy and those managed with physician-directed therapy.

Large, randomized, prospective clinical studies comparing outcomes, including OS and disease-specific survival, quality of life, and adverse events, between assay-directed therapy and physician-directed therapy, with outcomes are needed.

Chemosensitivity Assays

Chemosensitivity assays are designed to select the most appropriate chemotherapy regimens for a given tumor type, and would therefore ideally be associated with high positive predictive values (PPVs) for clinical response. The critical type of evidence needed to establish the effectiveness of chemosensitivity assays would come from comparative studies of assay-guided therapy versus physician-directed therapy. Relevant outcomes would include OS and disease-specific survival, as well as quality of life and adverse events.

The 2002 TEC Assessment (10) and 2004 systematic review (11) identified 9 comparative studies, two of which were randomized. (26-34) These authors reported that significant advantages for assay-guided therapy in terms of tumor response did not translate into survival differences. Response rate differences seen in other nonrandomized comparative studies may be attributable to bias or confounding, and survival outcomes were rarely reported.

Comparative Studies Testing Outcome with Assay-Directed Therapy versus Physician-Chosen Therapy

In a case-control study, Moon et al retrospectively compared adenosine triphosphate (ATP) assay-based guided chemotherapy with empirical chemotherapy in unresectable non-small-cell lung cancer. (35) All of the patients who received ATP-assay-guided platinum-based doublet chemotherapy as first-line therapy received platinum-based chemotherapy combined with a nonplatinum drug, regardless of their in vitro platinum sensitivity; 14 patients had platinum-sensitive disease and 13 were platinum-resistant. Ninety-three matched controls (matched for performance status, stage, and chemotherapy regimen) were selected from a retrospective review of a database. In the empirical group, a nonplatinum drug was chosen, depending on physicians' discretion, along with a platinum agent determined by renal function and performance status. The primary endpoint was clinical response rate, assessed every 2 cycles of chemotherapy by the Response Evaluation Criteria in Solid Tumors (RECIST) criteria. The secondary endpoints were PFS and OS. The response rate and survival in both groups were not statistically different. The platinum-sensitive subgroup by ATP assay showed a higher response rate than the empirical group (71% vs. 38%, respectively; $p=0.02$), but there was no statistical significance between PFS or OS.

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	9 of 19

In a small nonrandomized comparative study (n=64), Iwahashi et al (36) reported on outcomes of chemosensitivity-guided chemotherapy compared to standard chemotherapy and no chemotherapy in patients with advanced gastric cancer. In some subsets, survival was improved in the CSC subgroup. However, given the small sample, additional studies are needed to confirm these findings and to extend them to other malignancies.

Cree et al (37) reported on a prospective, randomized trial of chemosensitivity assay-directed chemotherapy versus physician's choice in patients with recurrent platinum-resistant ovarian cancer. The primary aim of this randomized trial was to determine response rate and PFS following chemotherapy in patients who had been treated according to an ATP-based tumor chemosensitivity assay in comparison with the physician's choice. A total of 180 patients were randomized to assay-directed therapy (n=94) or physician-choice chemotherapy (n=86). Median follow-up at analysis was 18 months; response was assessable in 147 (82%) patients: 31.5% achieved a partial or complete response in the physician-choice group compared with 40.5% in the assay-directed group (26% vs. 31% by intention-to-treat [ITT] analysis, respectively). ITT analysis showed a median PFS of 93 days in the physician's-choice group and 104 days in the assay-directed group (hazard ratio [HR]= 0.8, NS). No difference was seen in OS between the groups, although 12 of 39 patients (41%) who crossed over from the physician's-choice arm obtained a response. Increased use of combination therapy was seen in the physician's-choice arm during the study as a result of the observed effects of assay-directed therapy in patients. The authors concluded that this small RCT documented a trend toward improved response and PFS for assay-directed treatment and that chemosensitivity testing might provide useful information in some patients with ovarian cancer. They also noted that the ATP-based tumor chemosensitivity assay remains an investigational method in this condition.

Correlational Studies

Prospective. Kim et al reported the results of a prospective, multicenter clinical trial designed to define the accuracy of the ATP-based chemotherapy response assay in gastric cancer patients receiving paclitaxel and cisplatin chemotherapy, by comparing clinical response and the ATP-assay results. (38) The primary endpoint of the study was to assess accuracy of the ATP-assay results, and the secondary endpoint was to find the best method of defining in vitro chemosensitivity. Forty-eight patients with chemotherapy naïve locally advanced or metastatic gastric cancer were treated with combination chemotherapy after a tissue specimen was obtained for the ATP assay. Tumor response was assessed by World Health Organization (WHO) criteria using a computed tomography (CT) scan after every 2 cycles of chemotherapy. Both laboratory technicians and physicians were blinded to the assay or clinical results. Thirty-six patients were evaluable for both in vitro and in vivo responses. Using a chemosensitivity index method, the specificity of the ATP assay was 95.7% (95% CI, 77.2 to 99.9%), sensitivity 46.2% (95% CI, 19.2 to 74.9%), PPV was 85.7% (95% CI, 42.1 to 99.6%) and NPV was 75.9% (95% CI, 55.1 to 89.3%). Median PFS was 4.2 months (95% CI, 3.4 to 5.0) and median OS was 11.8 months (95% CI, 9.7 to 13.8). The in vitro chemosensitive group showed a higher response rate (85.7% vs 24.1%, respectively; p=0.005) compared to the chemoresistant group. The authors concluded that the ATP assay could predict clinical response to paclitaxel and cisplatin chemotherapy with high accuracy in advanced gastric cancer and that the study supported the use of the ATP assay in further validation studies.

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	10 of 19

In a European study, Ugurel et al reported on a nonrandomized, prospective, Phase 2 study of 53 evaluable patients with metastatic melanoma. (39) All 53 received assay-directed therapy. This study found a 36% response rate in patients with chemosensitive tumors compared with 16% in those with chemoresistant tumors. Based on these preliminary results, a Phase 3 study is to follow.

Rutherford et al reported results from a prospective, noninterventional, multicenter cohort study that was designed to assess whether the ChemoFX assay was predictive of outcomes among women with histologically confirmed epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer.(40) Three hundred thirty five patients were enrolled and treated with one of 15 study protocols, with treating physicians blinded to the ChemoFX assay result. Two hundred sixty-two patients (78.2% of total) had both available clinical follow up data and a ChemoFX result. Cancer cells were classified based on the ChemoFX result as sensitive, intermediate, or resistant to each of several chemotherapeutic agents. Patients treated with an assay-sensitive regimen had a PFS of median 8.8 months, compared with 5.9 months for those with assay-intermediate or –resistant regimens (HR 0.67, p=0.009). Mean overall survival was 37.5 months for patients treated with an assay-sensitive regimen, compared with 23.9 months for those with assay-intermediate or –resistant regimens (HR 0.67, P=0.010).

Rutherford et al reported results from a prospective, noninterventional, multicenter cohort study that was designed to assess whether the ChemoFX assay was predictive of outcomes among women with histologically confirmed epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. (40) Three hundred thirty-five patients were enrolled and treated with 1 of 15 study protocols, with treating physicians blinded to the ChemoFX assay result. Two hundred sixty-two patients (78.2% of total) had both available clinical follow-up data and a ChemoFX result. Cancer cells were classified based on the ChemoFX result as sensitive, intermediate, or resistant to each of several chemotherapeutic agents. Patients treated with an assay-sensitive regimen had a PFS of median 8.8 months, compared with 5.9 months for those with assay-intermediate or -resistant regimens (HR=0.67, p=0.009). Mean overall survival was 37.5 months for patients treated with an assay-sensitive regimen, compared with 23.9 months for those with assay-intermediate or -resistant regimens (HR=0.67, p=0.010).

In a follow-up analysis, Tian et al evaluated the ChemoFX's ability to predict PFS by comparing the association when the assayed therapy matched the administered therapy (match) with the association when the assayed therapy was randomly selected (mismatch). (41) The authors generated a simulation in which the average prognostic value of assay results for multiple different therapies was generated using the assay results for mismatch, in which the assay result for one treatment was randomly selected from the (up to) 15 designated therapies with equal probability for each patient. Based on 3000 repeated simulated resamplings, the mean HR for cases of mismatch was 0.81 (95% range, 0.66 to 0.99), which the authors suggest indicates that patients with a mismatch had less benefit when treated with an assay-sensitive therapy.

Strengths of this study include its prospective design with physicians blinded to the assay results, which reduces the risk of bias in patient selection or measurement of outcomes. However, because the selection of chemotherapeutic agent was, by design, not influenced by the ChemoFX assay, the impact on health outcomes cannot be determined.

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	11 of 19

Krivak et al reported results from a subsequent prospective, observational, multicenter study to determine whether sensitivity to carboplatin and/or paclitaxel is associated with disease progression among patients with primary epithelial ovarian cancer following initial treatment with a platinum/taxane regimen. (42) A total of 462 patients were enrolled, with 276 evaluable for inclusion in the analysis. Assay results for carboplatin and paclitaxel were available for 231 and 226 patients, respectively, with 44 (19.1%) patients identified as carboplatin-resistant and 49 (21.7%) identified as paclitaxel resistant. Carboplatin-resistant patients were at a higher risk of disease progression compared with nonresistant patients (HR=1.87; 95% CI, 1.29 to 2.70; $p<0.001$).

In a similar study design, Salom et al conducted a prospective, noninterventional, multicenter cohort study to assess whether the Microculture Kinetic (MiCK) assay (now called CorrectChemo) was predictive of outcomes among women with epithelial ovarian cancer.(43) Data from 150 women with any stage of cancer with specimens suitable for MiCK assay were included. Chemosensitivity was expressed as kinetic units following each dose of drug in the MiCK assay and reported as mean, minimum, and maximum. For each patient, the “best” chemotherapy was defined as any single drug or combination of drugs in the patient’s MiCK assay that had the highest kinetic units. Patients’ regimens were at the discretion of their treating physicians, who were blinded to the MiCK assay results. Overall survival Stage III or IV disease was longer if patients received a chemotherapy which was considered “best” by the MiCK assay, compared to shorter survival in patients who received a chemotherapy that was not the best. (HR 0.23, $P < 0.01$).

Jung et al conducted a single-center prospective study to determine whether sensitivity to paclitaxel and carboplatin, determined using the Histoculture Drug Resistance Assay (HDRA), was predictive of outcomes among women with advanced epithelial ovarian cancer.(44) The study included 104 patients with epithelial ovarian cancer, all of whom had undergone initial surgery and were treated with paclitaxel and carboplatin therapy. Tumor cells’ sensitivity to the chemotherapy agents was classified as sensitive, intermediate, or resistant to paclitaxel, carboplatin, or both, based on the HDRA. Patients whose tumors were sensitive to both drugs had a lower recurrence rate than those who had resistance to both drugs (29.2% vs 69.8%, $p=0.02$) and had a longer PFS (35 months vs 16 months, $p=0.025$).

While these studies establish that the results of chemosensitivity assays are correlated with outcome, they do not evaluate how the test may alter clinical decision-making and whether changes in management based on the test improve outcomes.

Retrospective: A number of retrospective studies have evaluated the association with various chemosensitivity assays and clinical outcomes in several tumor types, most commonly epithelial ovarian cancer. Some representative studies are discussed next.

Gallion et al conducted a retrospective study (45) that evaluated the association of ChemoFX® test results with the treatment response of 256 patients with ovarian or peritoneal cancer who had been treated with at least one cycle of postsurgical chemotherapy. A subset of 135 patients had an exact match between drugs assayed and received; the rest had only a partial match. Predictive values were not reported nor were they calculable. For the subset of 135, in a multivariable analysis, ChemoFX® was an independent significant predictor ($p=0.006$) of PFS along with 2 other clinical variables. HR for resistant versus sensitive was 2.9 (95% CI, 1.4 to 6.30) and was 1.7 (95% CI: 1.2 to 2.5) for resistant

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	12 of 19

versus intermediate. The median PFS was 9 months for the resistant group, 14 months for the intermediate group, and had not been achieved for the sensitive group.

Herzog et al included 147 patients from the above study by Gallion et al. (45) and reported on a total of 192 women with advanced-stage primary ovarian cancer, of whom 175 had tumors that were tested for in vitro chemosensitivity to platinum therapy using ChemoFX. (46) Tumors were classified as responsive, intermediately responsive, or nonresponsive to chemotherapy. Seventy-eight percent were categorized as responsive or intermediately responsive, and 22% were nonresponsive. Median OS was 72.5 months for patients with tumors categorized as responsive, 48.6 months for intermediately responsive, and 28.2 months for nonresponsive ($p=0.03$; HR 0.70; 95% CI, 0.50 to 0.97). The authors concluded that the result of chemosensitivity testing with a drug response marker for therapy was predictive of OS in patients with primary ovarian cancer.

In a smaller study, Grigsby et al conducted a retrospective analysis to assess the association of pretreatment chemosensitivity to cisplatin with clinical outcomes among 33 women with cervical cancer.(47) Tumor cell sensitivity to cisplatin was categorized as responsive, intermediately responsive, or nonresponsive with the ChemoFX assay. Patients with responsive or intermediately responsive tumors had a 2-year recurrence free survival of 87%, compared to 58% for those with nonresponsive tumors ($p=0.036$).

Lee et al conducted a retrospective study of the histoculture drug response assays (HDRA) assay in 79 patients with ovarian cancer. (48) Tissue samples were assessed for 11 chemotherapeutic agents and found the highest inhibition rates in carboplatin (49.2%), topotecan (44.7%), and belotecan (39.7%). These inhibition rates were higher than in cisplatin (34.7%), the traditional drug used to treat epithelial ovarian cancer. A subset of 37 patients with FIGO stage II/IV stage III or IV epithelial ovarian serous adenocarcinoma who had been treated with at least 3 cycles of carboplatin chemotherapy was assessed to compare outcomes between carboplatin-sensitive and -resistant patients. Multiple comparison and regression analyses established a cut-off value of 40% inhibition rate in response to 50 ug/mL carboplatin to determine sensitivity or resistance. This selected cut-off had a disease-free survival of 23.2 months (95% CI: 6.3-55.3) and 13.8 months (95% CI: 4.9-35.6) in the carboplatin-sensitive and carboplatin-resistant groups respectively ($p<0.05$). Overall survival between the 2 groups did not differ significantly, with carboplatin-sensitive patients having a mean 60.4 months and carboplatin-resistant patients having 37.3 months ($p=0.621$).

Strickland et al conducted a retrospective evaluation of the association between chemosensitivity to anthracyclines, measured by the drug-induced apoptosis MiCK assay (now called the Correct Chemo assay), among 109 patients with adult-onset acute myelogenous leukemia.(49) Patients were treated with a "7 plus 3" chemotherapy regimen. Chemosensitivity was expressed as maximal kinetic units following each dose of drug in the MiCK assay. Receiver-operator characteristic curve analysis and logistic regression were used to determine the optimal cutoff for chemosensitivity response to discriminate between chemoresponder and non-responder. Patients determined to be chemoresponders to idarubicin were more likely to have complete response to chemotherapy (72%) than those who were non-responders ($p=0.01$). Data for the patient cohort were collected over a 14 year period from 1996-2010, which may limit the generalizability of the results to currently-used

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	13 of 19

chemotherapy regimens. In addition, the MiCK assay is limited by lack of standardized cutoffs to discriminate responders from nonresponders.

Other retrospective studies have evaluated the association between chemosensitivity as measured by other assay types. Von Heideman et al evaluated the semi-automated fluorometric microculture cytotoxicity assay (FMCA) in 112 patients (125 samples) with ovarian cancer and concluded that samples from patients with clinical response were more sensitive to most drugs than samples from non-responding patients. (50)

Section Summary:

The most direct evidence related to the effectiveness of chemosensitivity assays in the management of patients with cancer comes from several studies which compare outcomes for patients managed with an ATP-based tumor chemosensitivity assay with those managed with standard care, including 1 randomized controlled trial. Although some improvements in tumor response were noted, no differences between OS or PFS were seen. A number of retrospective and prospective studies of several different chemosensitivity assays, including the ATP-based tumor chemosensitivity assay, the CorrectChemo assay, and the ChemoFX assay, suggest that patients whose tumors have higher chemosensitivity have better outcomes. However, additional studies to determine whether the clinical use of in vitro chemosensitivity testing leads to better outcomes are needed.

Ongoing and Unpublished Clinical Trials

Table 1: Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing NCT02137811 ^a	Chart Review of Demographics and Clinical Outcomes and Physician Attitudes in Patients Who Have Received the Microculture Kinetic (MiCK) Apoptosis Test	400	Jan 2017

NCT: national clinical trial

^a Denotes industry-sponsored or cosponsored trial

Practice Guidelines and Position Statements

National Comprehensive Cancer Network (NCCN)

The 2015 NCCN Guidelines for the treatment of epithelial ovarian cancer, fallopian tube cancer, and primary peritoneal cancer (v 1.2015) states the following, "Chemosensitivity/resistance and/or other biomarker assays are being used in some NCCN Member Institutions. The current level of evidence is not sufficient to supplant standard-of-care chemotherapy (category 3)." (51)

The American Society of Clinical Oncology Clinical Practice Guideline Update on the Use of Chemotherapy Sensitivity and Resistance Assays, 2011 also does not recommend use of chemotherapy sensitivity and resistance assays, unless in a clinical trial setting. (52)

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	14 of 19

U.S. Preventive Services Task Force Recommendations

Not applicable

Summary

There are only a few comparative studies that evaluate use of a chemosensitivity assay to select chemotherapy versus standard care, and these studies do not report significant differences in outcomes between groups. A larger number of studies have used correlational designs that evaluate the association between assay results and already known patient outcomes. These studies report that results of chemosensitivity and chemoresistance assays are predictive of outcomes. However, these studies do not evaluate whether these assays lead changes in management, and whether any changes in management lead to improved outcomes. In addition, interpretation of these studies is limited by heterogeneity in test methodology, tumor type, patient population, and chemotherapeutic. As a result, the clinical utility of chemoresistance and chemosensitivity assays has not been determined, and data are insufficient to determine whether use of the test to select chemotherapy regimens for individual patients will improve outcomes. Therefore, this testing is considered **not medically necessary**.

Medicare National Coverage

Not applicable

References

1. Bird MC, Godwin VA, Antrobus JH, et al. Comparison of in vitro drug sensitivity by the differential staining cytotoxicity (DiSC) and colony-forming assays. *Br J Cancer*. Apr 1987; 55(4):429-431. PMID 3580265
2. Nagourney RA, Blitzer JB, Shuman RL, et al. Functional profiling to select chemotherapy in untreated, advanced or metastatic non-small cell lung cancer. *Anticancer Res*. Oct 2012; 32(10):4453-4460. PMID 23060572
3. Nagourney RA. Ex vivo programmed cell death and the prediction of response to chemotherapy. *Curr Treat Options Oncol*. Mar 2006; 7(2):103-110. PMID 16455021
4. Csoka K, Larsson R, Tholander B, et al. Cytotoxic drug sensitivity testing of tumor cells from patients with ovarian carcinoma using the fluorometric microculture cytotoxicity assay (FMCA). *Gynecol Oncol*. Aug 1994; 54(2):163-170. PMID 7520407
5. Yung WK. In vitro chemosensitivity testing and its clinical application in human gliomas. *Neurosurg Rev*. 1989; 12(3):197-203. PMID 2682352
6. Kern DH, Weisenthal LM. Highly specific prediction of antineoplastic drug resistance with an in vitro assay using suprapharmacologic drug exposures. *J Natl Cancer Inst*. Apr 4 1990; 82(7):582-588. PMID 2313735
7. Anticancer I. HISTOCULTURE DRUG RESPONSE ASSAY - HDRA. Available online at: http://www.anticancer.com/HDRA_ref.html. Last accessed May, 2015.
8. Precision Therapeutics Incorporated. ChemoFx. Available online at: <http://www.chemofx.com/index.html>. Last accessed May, 2015.

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	15 of 19

9. DiaTech Oncology. MiCK Assay. Available online at: <http://correctchemo.com/>. Last accessed May, 2015.
10. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Chemotherapy sensitivity and resistance assays. TEC Assessments 2002; Volume 17, Tab 12. PMID 12166470
11. Samson DJ, Seidenfeld J, Ziegler K, et al. Chemotherapy sensitivity and resistance assays: a systematic review. *J Clin Oncol*. Sep 1 2004; 22(17):3618-3630. PMID 15289487
12. Brown E, Markman M. Tumor chemosensitivity and chemoresistance assays. *Cancer* Mar 15 1996; 77(6):1020-1025. PMID 8635118
13. Eltabbakh GH. Extreme drug resistance assay and response to chemotherapy in patients with primary peritoneal carcinoma. *J Surg Oncol*. Mar 2000; 73(3):148-152. PMID 10738268
14. Eltabbakh GH, Piver MS, Hempling RE, et al. Correlation between extreme drug resistance assay and response to primary paclitaxel and cisplatin in patients with epithelial ovarian cancer. *Gynecol Oncol*. Sep 1998; 70(3):392-397. PMID 9790793
15. Mehta RS, Bornstein R, Yu IR. Breast cancer survival and in vitro tumor response in the extreme drug resistance assay. *Breast Cancer Res Treat*. Apr 2001; 66(3):225-237. PMID 11510694
16. Holloway RW, Mehta RS, Finkler NJ, et al. Association between in vitro platinum resistance in the EDR assay and clinical outcomes for ovarian cancer patients. *Gynecol Oncol*. Oct 2002; 87(1):8-16. PMID 12468336
17. Ellis RJ, Fabian CJ, Kimler BF, et al. Factors associated with success of the extreme drug resistance assay in primary breast cancer specimens. *Breast Cancer Res Treat*. Jan 2002; 71(2):95-102. PMID 11881914
18. Loizzi V, Chan JK, Osann K, et al. Survival outcomes in patients with recurrent ovarian cancer who were treated with chemoresistance assay-guided chemotherapy. *Am J Obstet Gynecol*. Nov 2003; 189(5):1301-1307. PMID 14634558
19. Tiersten AD, Moon J, Smith HO, et al. Chemotherapy resistance as a predictor of progression-free survival in ovarian cancer patients treated with neoadjuvant chemotherapy and surgical cytoreduction followed by intraperitoneal chemotherapy: a Southwest Oncology Group Study. *Oncology*. 2009; 77(6):395-399. PMID 20130422
20. Matsuo K, Eno ML, Im DD, et al. Clinical relevance of extent of extreme drug resistance in epithelial ovarian carcinoma. *Gynecol Oncol*. Jan 2010; 116(1):61-65. PMID 19840886
21. Matsuo K, Bond VK, Eno ML, et al. Low drug resistance to both platinum and taxane chemotherapy on an in vitro drug resistance assay predicts improved survival in patients with advanced epithelial ovarian, fallopian and peritoneal cancer. *Int J Cancer*. Dec 1 2009; 125(11):2721-2727. PMID 19530239
22. Matsuo K, Bond VK, Im DD, et al. Prediction of chemotherapy response with platinum and taxane in the advanced stage of ovarian and uterine carcinosarcoma: a clinical implication of in vitro drug resistance assay. *Am J Clin Oncol*. Aug 2010; 33(4):358-363. PMID 19875949

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	16 of 19

23. Matsuo K, Eno ML, Im DD, et al. Chemotherapy time interval and development of platinum and taxane resistance in ovarian, fallopian, and peritoneal carcinomas. *Arch Gynecol Obstet.* Feb 2010; 281(2):325-328. PMID 19455347
24. Karam AK, Chiang JW, Fung E, et al. Extreme drug resistance assay results do not influence survival in women with epithelial ovarian cancer. *Gynecol Oncol.* Aug 2009; 114(2):246-252. PMID 19500821
25. Hetland TE, Kaern J, Skrede M, et al. Predicting platinum resistance in primary advanced ovarian cancer patients with an in vitro resistance index. *Cancer Chemother Pharmacol.* May 2012; 69(5):1307-1314. PMID 22302409
26. Cortazar P, Gazdar AF, Woods E, et al. Survival of patients with limited-stage small cell lung cancer treated with individualized chemotherapy selected by in vitro drug sensitivity testing. *Clin Cancer Res.* May 1997; 3(5):741-747. PMID 9815744
27. Gazdar AF, Steinberg SM, Russell EK, et al. Correlation of in vitro drug-sensitivity testing results with response to chemotherapy and survival in extensive-stage small cell lung cancer: a prospective clinical trial. *J Natl Cancer Inst.* Jan 17 1990; 82(2):117-124. PMID 2152944
28. Kurbacher CM, Cree IA, Bruckner HW, et al. Use of an ex vivo ATP luminescence assay to direct chemotherapy for recurrent ovarian cancer. *Anticancer Drugs.* Jan 1998; 9(1):51-57. PMID 9491792
29. Shaw GL, Gazdar AF, Phelps R, et al. Individualized chemotherapy for patients with non-small cell lung cancer determined by prospective identification of neuroendocrine markers and in vitro drug sensitivity testing. *Cancer Res.* Nov 1 1993; 53(21):5181-5187. PMID 8221655
30. Shaw GL, Gazdar AF, Phelps R, et al. Correlation of in vitro drug sensitivity testing results with response to chemotherapy and survival: Comparison of non-small cell lung cancer and small cell lung cancer. *J Cell Biochem Suppl.* 1996; 24:173-185. PMID 8806100
31. Von Hoff DD, Kronmal R, Salmon SE, et al. A Southwest Oncology Group study on the use of a human tumor cloning assay for predicting response in patients with ovarian cancer. *Cancer.* Jan 1 1991; 67(1):20-27. PMID 1985717
32. Von Hoff DD, Sandbach JF, Clark GM, et al. Selection of cancer chemotherapy for a patient by an in vitro assay versus a clinician. *J Natl Cancer Inst.* Jan 17 1990; 82(2):110-116. PMID 2403593
33. Wilbur DW, Camacho ES, Hilliard DA, et al. Chemotherapy of non-small cell lung carcinoma guided by an in vitro drug resistance assay measuring total tumour cell kill. *Br J Cancer.* Jan 1992; 65(1):27-32. PMID 1310250
34. Xu JM, Song ST, Tang ZM, et al. Predictive chemotherapy of advanced breast cancer directed by MTT assay in vitro. *Breast Cancer Res Treat.* Jan 1999; 53(1):77-85. PMID 10206075
35. Moon YW, Sohn JH, Kim YT, et al. Adenosine triphosphate-based chemotherapy response assay (ATP-CRA)-guided versus empirical chemotherapy in unresectable non-small cell lung cancer. *Anticancer Res.* Oct 2009; 29(10):4243-4249. PMID 19846981
36. Iwahashi M, Nakamori M, Nakamura M, et al. Individualized adjuvant chemotherapy guided by chemosensitivity test sequential to extended surgery for advanced gastric cancer. *Anticancer Res.* Sep-Oct 2005; 25(5):3453-3459. PMID 16101163

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	17 of 19

37. Cree IA, Kurbacher CM, Lamont A, et al. A prospective randomized controlled trial of tumour chemosensitivity assay directed chemotherapy versus physician's choice in patients with recurrent platinum-resistant ovarian cancer. *Anticancer Drugs*. Oct 2007; 18(9):1093-1101. PMID 17704660
38. Kim JH, Lee KW, Kim YH, et al. Individualized tumor response testing for prediction of response to Paclitaxel and Cisplatin chemotherapy in patients with advanced gastric cancer. *J Korean Med Sci*. May 2010; 25(5):684-690. PMID 20436702
39. Ugurel S, Schadendorf D, Pfohler C, et al. In vitro drug sensitivity predicts response and survival after individualized sensitivity-directed chemotherapy in metastatic melanoma: a multicenter phase II trial of the Dermatologic Cooperative Oncology Group. *Clin Cancer Res*. Sep 15 2006; 12(18):5454-5463. PMID 17000680
40. Rutherford T, Orr J, Jr., Grendys E, Jr., et al. A prospective study evaluating the clinical relevance of a chemoresponse assay for treatment of patients with persistent or recurrent ovarian cancer. *Gynecol Oncol*. Nov 2013; 131(2):362-367. PMID 23954900
41. Tian C, Sargent DJ, Krivak TC, et al. Evaluation of a chemoresponse assay as a predictive marker in the treatment of recurrent ovarian cancer: further analysis of a prospective study. *Br J Cancer*. Aug 26 2014; 111(5):843-850. PMID 25003664
42. Krivak TC, Lele S, Richard S, et al. A chemoresponse assay for prediction of platinum resistance in primary ovarian cancer. *Am J Obstet Gynecol*. Jul 2014;211(1): 68 e61-68. PMID 24530815
43. Salom E, Penalver M, Homesley H, et al. Correlation of pretreatment drug induced apoptosis in ovarian cancer cells with patient survival and clinical response. *J Transl Med*. 2012;10:162. PMID 22873358
44. Jung PS, Kim DY, Kim MB, et al. Progression-free survival is accurately predicted in patients treated with chemotherapy for epithelial ovarian cancer by the histoculture drug response assay in a prospective correlative clinical trial at a single institution. *Anticancer Res*. Mar 2013;33(3):1029-1034. PMID 23482777
45. Gallion H, Christopherson WA, Coleman RL, et al. Progression-free interval in ovarian cancer and predictive value of an ex vivo chemoresponse assay. *Int J Gynecol Cancer*. Jan-Feb 2006; 16(1):194-201. PMID 16445633
46. Herzog TJ, Krivak TC, Fader AN, et al. Chemosensitivity testing with ChemoFx and overall survival in primary ovarian cancer. *Am J Obstet Gynecol*. Jul 2010; 203(1):68 e61-66. PMID 20227055
47. Grigsby PW, Zigelboim I, Powell MA, et al. In vitro chemoresponse to cisplatin and outcomes in cervical cancer. *Gynecol Oncol*. Jul 2013; 130(1):188-191. PMID 23583416
48. Lee JH, Um JW, Lee JH, et al. Can immunohistochemistry of multidrug-resistant proteins replace the histoculture drug response assay in colorectal adenocarcinomas? *Hepatogastroenterology*. Jun 2012; 59(116):1075-1078. PMID 22580657
49. Strickland SA, Raptis A, Hallquist A, et al. Correlation of the microculture-kinetic drug-induced apoptosis assay with patient outcomes in initial treatment of adult acute myelocytic leukemia. *Leuk Lymphoma*. Mar 2013; 54(3):528-534. PMID 22924433

Section:	Medicine	Effective Date:	July 15, 2015
Subsection:	Oncology	Original Policy Date:	December 7, 2011
Subject:	In Vitro Chemoresistance and Chemosensitivity Assays	Page:	18 of 19

50. von Heideman A, Tholander B, Grundmark B, et al. Chemotherapeutic drug sensitivity of primary cultures of epithelial ovarian cancer cells from patients in relation to tumour characteristics and therapeutic outcome. *Acta Oncol.* Feb 2014;53(2):242-250. PMID 23713890
51. National Comprehensive Cancer Network (NCCN). Guidelines v1.2015 Ovarian Cancer. 2015. Available online at: http://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf. Last accessed May, 2015.
52. Burstein HJ, Mangu PB, Somerfield MR, et al. American Society of Clinical Oncology Clinical Practice Guideline Update on the Use of Chemotherapy Sensitivity and Resistance Assays. *J Clin Oncol.* Aug 20 2011; 29(24):3328-3330. PMID 21788567

Policy History

Date	Action	Reason
December 2011	New Policy	
June 2012	Update Policy	Policy statement changed to not medically necessary.
June 2013	Update Policy	Policy updated with literature review. References 2,3, 21, and 40 added, some reordered. No change to policy statement.
June 2014	Update Policy	Policy updated with literature review. References 6-8, 40-42, 45, and 47- 48 added. Background and rationale reorganized. No changes to policy statements.
June 2015	Update Policy	Policy updated with literature review. References 4, 41-42, and 50 added. "ChemoFx" and "CorrectChemo" added to the list on investigational chemosensitivity assays; policy statements otherwise unchanged.

Keywords

AntiCancer, Inc; Histoculture Drug Response Assay
 Chemoresistance Assay
 Chemosensitivity Assay
 Cytoprint
 EDR Assay
 Extreme Drug Resistance Assay
 Histoculture Drug Response Assay
 Oncotech, EDR Assay

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on June 19, 2015 and is effective July 15, 2015.

Signature on File

Deborah M. Smith, MD, MPH